





INTRODUCTION

The systematic mapping of lava flows in the Tharsis region has been compiled into a series of 16 maps at 1:2,000,000 scale. This work provides information on the sources and areal extent of the lava flows, on their eruptive sequences and relative ages, and on relations between the flows and geologic structure in the largest, most active tectonic and volcanic province on Mars. Some of the maps were made from controlled Viking photomosaics published as quarter quadrangles in the Atlas of Mars Topographic Series (U.S. Geological Survey, 1979) and tied to the Viking control net. Where these photomosaics were not available, larger scale analog photomosaics tied to the Mariner 9 control net were used. These maps were subsequently reduced to the 1:2,000,000 scale, but slight discrepancies may occur in places between features referred to coordinates on the two types of bases.

Mariner 9 orbital images of the region show a few major flow units, mapped around Olympus Mons by Carr (1975) and by Morris and Dwornik (1978), around Arisa Mons by Maursky, Dial and Strobel (1978), and around Alba Patera by Wise (1979). However, flow lobes characteristic of individual lava extrusions were difficult to recognize on the wide-angle camera frames used for geologic mapping. With the acquisition of high-resolution images by the Viking mission, numerous individual lava flows in the Tharsis region were identified and mapped in detail over large areas (Schaber and others, 1978; Scott and others, 1979). Although the geologic investigation was mostly directed toward the mapping of lava flows and the determination of their eruptive sequences, structural features such as faults, fractures and the basal scarp around Olympus Mons were also mapped and dated relative to the flow units. In this way a sequence of tectonic episodes was determined in conjunction with the major volcanic events.

The Tharsis volcanic province as defined in this study covers some 18 million square kilometers. It is approximately rectangular, extending from lat 40° S to lat 45° N between long 90° and long 125° (fig. 1). Within this province occur the four largest and youngest volcanoes on Mars: Olympus Mons, Arisa Mons, Pavonis Mons, and Ascraeus Mons; the latter three collectively form the large elevated area named the Tharsis Montes. Other major physiographic and structural features are Alba Patera, an ancient low-relief volcano of great size; Achernis Fossae, thought to be a volcano-tectonic structure, possibly similar to Alba Patera, but older; and Syria Planum, a very large domical uplift southeast of the Tharsis Montes.

The relative ages of major eruptive sequences were determined mainly by their stratigraphic relations and by morphology of the flows. Crater counts on the various units were made to verify these age relations and to obtain some degree of correlation between flows in widely separated areas where overlap relations could not be established. Crater size-frequency distributions were calculated from crater counts on moderate-resolution (180-280 m/pixel) frames. In general, counts were made only within units having boundaries defined by standard photogeologic techniques. Craters with subdued or modified morphologies possibly indicative of preflow origin were not counted. Sources of error in crater counts included variations in cloud cover, sun-elevation angle, and resolution in images of different scales. Statistically valid data were obtained by counting craters within large areas of individual flows or geologic units; these areas range from about 3,600 km² to 176,500 km². All units have been assigned to the martian time-stratigraphic systems shown on the 1:25,000,000-scale map of Mars (Scott and Carr, 1978). Differences between these assignments and those on the small-scale map of the planet reflect revisions introduced by the detailed mapping.

GEOLOGIC SUMMARY

Martian lava flows are similar in morphology to those on Earth and the Moon. They commonly exhibit overlapping, lobate, and crenulated margins and occur chiefly as sheet flows or as channel- and tubed flows (Carr and others, 1977). Sheet flows are more common on the plains and on the lower, more gentle slopes of volcanoes. Their surfaces appear flat and smooth at moderate resolution, but at high resolution they exhibit concentric ridge-and-groove patterns subparallel to flow margins. Channel- and tube flows are more prevalent in the steeper slopes around volcanoes such as Olympus Mons and Arisa Mons, but also occur on relatively low-relief surfaces at Alba Patera and Ceratunus Fossae. Younger flows have rougher textures than older ones that have been smoothed by erosion and mantled to various degrees by collan deposits. The surface flows, like those on Earth, originated from the central vents of volcanoes or from radial fissures, or from fissure vents, and are generally far removed from the volcanic edifices. Of the 24 major lava-flow sequences mapped in the Tharsis region of Mars, 13 emanated mostly from two large source volcanoes, Olympus Mons and Arisa Mons. The youngest recent flows were extruded from large fissures in the high plains east of Olympus Mons and from the summit areas of the Tharsis volcanoes. The oldest flows erupted from calderas and associated fissures in the Tharsis region, and in the high plains east of Olympus Mons (15° S, 100° W), respectively, a large shield volcano and a high-ventilation but low-relief dome of the planet's northern hemisphere.

Basement and Nonvolcanic Units

Basement rocks (unit HNht) are undivided. They consist of both rough and smooth, highly fractured terrain: billy and cratered material; and cratered plains and cratered plains materials that form a large part of the ancient martian highlands (Scott and Carr, 1978). They occur mostly as relatively large blocks emplaced and partly buried by various flows. Around the periphery of Olympus Mons, however, these older rocks may be exposed in the base-camp and as upflood blocks protruding above the lava flows in places. Some of this material may also represent segments of overlapping aureoles that formerly covered the present site of this volcano and Olympus Mons flows that predate the basal

scarp. However, these various units cannot be separately distinguished in such small areas, and most exposed parts of the scarp complex have been mapped as basement material.

Channel and flood-plain deposits and large accumulations of collan material have been mapped in a few places. They have been relatively dated stratigraphically and by crater counts with respect to the Tharsis lava flows and contribute information on climate in the evolutionary history of the region. Landlides are common around the large volcanoes in the Tharsis region. They occur along and below the basal scarp on the west side of Olympus Mons and on the northwest flanks of Arisa, Pavonis, and Ascraeus Montes. Although the slide material is undivided on the lava flow maps, it consists chiefly of two end members. 1) rough, blocky rockfalls and rockslides near the head of a dissected area that grade downslope into; 2) debris or mudflow deposits that form thin lobate tongues with many narrow ridges concentric to the outermost flow front. These deposits at Arisa Mons and Pavonis Mons are older than flows from the crestral areas of these volcanoes but overlap slightly older flows on their flanks.

Tharsis Lava Flows

The major flow units are grouped very generally into broad relative-age categories. The eruptive sequence appears to have been continuous throughout the volcanic history of the region. Crater densities cited below are the number of craters larger than 1 km in diameter normalized to an area of 1 million km².

Older Flows—Alba Patera, Syria Planum, and the Aureoles of Olympus Mons—The flows from Alba Patera and Syria Planum have the highest crater densities, ranging from about 1800 for the youngest to 3200 for the oldest flows at each location. These figures are roughly comparable to those of younger and older lunar maria at the Apollo 11 and 12 sites respectively (Neukam and Wise, 1976). Estimates of the absolute ages of these and other flows vary widely, however, because of inherent uncertainties in the models postulated for martian integrated flux curves used to establish correlations between crater frequency and geologic age (Neukam and Wise, 1976; Hartmann, 1977; Soderblom, 1977).

The younger flows from Alba Patera partly bury some of the radial and concentric fault systems that transect the older lavas. This burial is particularly noticeable within and around the caldera at the crest of the ancient volcano where the late and waning eruptive stages yielded decreasing volumes of lava. The older flows from Alba Patera embay the highly dissected terrain around the east edge of Achernis Fossae in the Ductia quadrangle (MC2SE). These lavas also have penetrated the arcuate central part of this volcano-tectonic structure where they may have been overlapped by very young lavas of the Olympus plains and possibly also by the oldest aureole unit surrounding Olympus Mons.

At Syria Planum in the Phoenicis Lacus quadrangle (MC1SE), the younger flows also occur near the crest of the structural dome. They probably issued from a partly buried calderalike depression or from fissures subsequently obscured by the numerous flows. These lavas also appear to have buried many of the faults and fractures associated with or preceding channel development along Nectis Labrynthus. The older flows of Syria Planum embay and partly bury the landforms and faults of Claritas Fossae but are transected by some of the more recent faults. No contacts can be clearly distinguished between the older and younger flow units at either Syria Planum or Alba Patera. The mapped boundaries are speculative and are based mostly on slight variations in morphology as well as crater densities. The aureoles of grooved and ridged terrain surrounding Olympus Mons are classified with the older group of flows. Their relative ages with respect to most other flow units in the Tharsis region can be only broadly determined. Stratigraphic relations clearly show that the aureoles are overlapped by the flows of Olympus plains and the postscarp extrusives of Olympus Mons. There is some evidence, but less certain, that the aureole deposits may be overlapped in places (MC8SW) by the Tharsis Montes flows. Craters are of little value as age determinants because the ridged and grooved surfaces of the aureoles promote rapid deterioration of crater forms by mass wasting; their crater densities are abnormally low, for example, compared with younger adjacent plains that embay the aureoles in places. Morphologically, the aureoles seem to be older than most flows in the Tharsis region. On the basis of stratigraphic evidence and morphologic relations, the aureole units of Olympus Mons are provisionally placed just above the Alba Patera and Syria Planum lavas in the eruptive sequence.

Flows of Intermediate Age—Tharsis Montes, Alba Patera, and Ceratunus Fossae—This group of lava flows has the greatest range in crater densities—from about 430 to 1970, overlapping slightly with those in the older category. Many of the flow units originated from Arisa Mons (fig. 1) and are exposed from the upper to the lower slopes of the volcano. These flows show the greatest diversity in surface characteristics as well as in relative ages. The lower, older flows around Arisa Mons are more subdued by erosion and collan mantles than the younger and generally higher units. It is not known if this condition applies at initial venting and remnant thermal venting and subsequent cratering. The most recent volcanic activity at Alba Patera includes the complex fracture systems of that region and subsequently buried many of the fissures from which they were extruded. They extend southwesterly from their source area across the lava flows from Alba Patera and embay part of the highland terrain

between Olympus Mons and Tharsis Montes. Some may have erupted from fissures in this highland region. They are overlapped, in turn, by younger flows from Tharsis Montes and from Olympus plains.

Younger Flows—Olympus Mons, Olympus Plains, and the Volcanoes of Tharsis Montes—Crater densities for this group of extrusives range from about 90 to 350. They originated from the flanks and summit of Olympus Mons, from fissures in the plains east of the volcano, and from the crestral areas of Arisa, Pavonis, and Ascraeus Montes. Stratigraphic relations are clearly defined between the flows of Olympus Mons and those of the adjacent plains. These plains partly enclose Olympus Mons on the east within a topographically low area between the volcano and its aureole deposits. To the west of Olympus Mons no evidence exists of well-defined young flow fronts. Postscarp flows in this area are either buried by collan material or are so deeply eroded as to be unrecognizable. Postscarp lava flows either did not occur on the west side of Olympus Mons or were of such small volume that they were unable to surmount the upturned edges of the scarp. Boundaries between pre- and postscarp flow units on Olympus Mons are not distinct. Crater counts indicate that two slightly different age groups may be present, but the validity of the data is limited by difficulties in selecting representative areas for each unit and errors inherent in distinguishing between small impact and indigenous craters on the flanks of a volcano. Evidence that precarap lava flows exist is shown in places by flows sharply truncated at the scarp and by exposures of more mature surfaces in windows along its raised edge. Alternatively, vertical displacements producing the basal scarp around Olympus Mons may have occurred episodically. Intermittent lava flows down the flanks of the volcano throughout the interval of scarp formation could result in the present configuration of pre- and postscarp flows in different places.


Flows from near the summits of Arisa Mons, Pavonis Mons, and Ascraeus Mons are among the youngest in the Tharsis region. They appear to be relatively thin, with smooth surfaces and indistinct flow fronts; their boundaries with other units are difficult to define. These lava flows originated from large fractures and fissures along northeast-southwest faults transecting the crestral areas of the Tharsis Montes volcanoes.

STRUCTURE

Tectonism, as expressed by fractures and faults, culminated before the bulk of the lava flows in the Tharsis region were extruded. It did not cease altogether, however, but continued with diminishing intensity into the period of the youngest flows (Scott and Tanaka, 1980). Early episodes of major faulting were responsible for the highly disrupted surfaces at Claritas, Achernis, Ceratunus, and Mennela Fossae. Faulting continued in these areas at lesser scales and also at Alba Patera, on the flanks of Arisa Mons, and on the flanks of Arisa Mons. The density of faults associated with individual flow units, like crater density, reflects their relative ages. A summary of tectonic episodes as they relate to major volcanic events is shown in the Correlation of Map Units.

REFERENCES

- Carr, M. H., 1975, Geologic map of the Tharsis quadrangle of Mars: U.S. Geological Survey Miscellaneous Investigations Series Map I-893.
Carr, M. H., Greeley, Ronald, Blais, K. R., Guest, J. E., and Murray, J. B., 1977, Some martian volcanic features as viewed from the Viking Orbiters: Journal of Geophysical Research, v. 82, no. 28, p. 3983-4015.
Hartmann, W. K., 1978, Martian cratering V: Toward an empirical martian chronology, and its implications. Geophysical Research Letters, v. 5, no. 6, p. 450-452.
Kieffer, H. H., Martin, T. Z., Peterfreund, A. R., and Jakosky, B. M., 1977, Thermal and albedo mapping of Mars during the Viking primary mission: Journal of Geophysical Research, v. 82, no. 28, p. 4249-4291.
Maursky, Harold, Dial, A. L., Jr., and Strobel, M. E., 1978, Geologic map of the Phoenicis Lacus quadrangle of Mars: U.S. Geological Survey Miscellaneous Investigations Series Map I-896.
Morris, E. C., and Dwornik, S. L., 1978, Geologic map of the Amazonis quadrangle of Mars: U.S. Geological Survey Miscellaneous Investigations Series Map I-1049.
Neukam, Gerhard, and Wise, D. U., 1976, Mars: A standard crater count and possible new time scale: Science, v. 194, no. 4272, p. 1381-1387.
Schaber, G. G., Hartman, K. C., and Dial, A. L., Jr., 1978, Lava flow materials in the Tharsis region of Mars: Lunar and Planetary Science Conference, 9th Proceedings, p. 3433-3458.
Scott, D. H., and Carr, M. H., 1978, Geologic map of Mars: U.S. Geological Survey Miscellaneous Investigations Series Map I-1083, 1:25,000,000 scale.
Scott, D. H., Schaber, G. G., Hartman, K. C., and Dial, A. L., Jr., 1979, Lava flows of Tharsis Montes: U.S. National Aeronautics and Space Administration Technical Memorandum TM-60339, Reports of Planetary Geology Program, 1978-1979, p. 237-238.
Scott, D. H., and Tanaka, K. L., 1980, Mars Tharsis region: Volcanic-tectonic events in the stratigraphic record: Lunar and Planetary Science Conference, 11th Proceedings, p. 2403-2421.
Soderblom, L. A., 1977, Historical variations in the density and distribution of impacting debris in the inner solar system: Evidence from planetary images in impact and explosion cratering, in Roddy, D. J., Pepin, R. O., and Merrill, R. B., eds., Impact and explosion cratering: New York, Pergamon Press, p. 629-633.
U.S. Geological Survey, 1979, and subsequent years, Atlas of Mars Topographic Series, Miscellaneous Investigations Series Maps, 1:2,000,000 scale.
Wise, D. U., 1979, Geologic map of the Arcadia quadrangle of Mars: U.S. Geological Survey Miscellaneous Investigations Series Map I-1154.



Crater populations on individual map units are indicated by the letter N, which refers to the number of craters counted larger than 1.0 km in diameter, normalized to an area of 1 million km².

DESCRIPTION OF MAP UNITS

YOUNGER FLOWS

OLYMPUS PLAINS FLOWS: Occur around basal scarp of Olympus Mons. Surfaces relatively smooth with many lobate scarps; crater density very low (N < 50). Overlap all adjacent units including flows from Olympus Mons (units Aam1, Aam2). In places elongate tongues are centered along fissure systems in plains east of Olympus Mons. *Interpretation:* Youngest lava flows in Tharsis region; extruded from faults and fissures in high plains east of Olympus Mons.

ARISA MONS FLOW UNIT 5: Forms smooth mantles on northeast and southwest upper slopes of Arisa Mons and partly fills caldera. Overlaps Tharsis Montes flows but boundaries between units poorly defined. Crater density very low (N < 100). *Interpretation:* Youngest lava flows from Arisa Mons originating from isolated vents near summit and within caldera.

ASCRAEUS MONS FLOWS: Similar in appearance and occurrence to youngest flow units from Arisa Mons and Pavonis Mons. *Interpretation:* Youngest flows from Ascraeus Mons.

PAVONIS MONS FLOW UNIT 2: Occurs on northeast and southwest sides of Pavonis Mons and inside caldera. Similar in appearance and occurrence to Arisa Mons flow unit 5. *Interpretation:* Youngest flows from Pavonis Mons.

PAVONIS MONS FLOW UNIT 1: Occurs on northwest side of Pavonis Mons. Lobate flow fronts visible in places. Overlaps Tharsis Montes flow unit but is overlapped by slide material. Crater density low (N = 100-200). *Interpretation:* Young lava flows extruded from central area and flanks of Olympus Mons after formation of basal scarp.

OLYMPUS MONS PRESCARP FLOWS: Small patches largely inferred along upturned edges of basal scarp in northwest and southeast side of Olympus Mons. Mapped only where lava flows extending down flanks of volcano are sharply truncated in places along scarp, or where older appearing surfaces occur as windows within younger flows (unit Aam2). *Interpretation:* Lava flows extruded before scarp formed at base of Olympus Mons.

THARSIS MONTES FLOWS: Form elongate lobes extending downslope from upper parts of Arisa, Pavonis, and Ascraeus Montes (collectively referred to as Tharsis Montes). Overlap and appear topographically sharper than other flows (units Aam1, Aam2, Aam3) exposed on lower flanks of Arisa Mons. Cut by low faults. Crater density range, N = 130-570. *Interpretation:* Late-stage lava flows extruded from Arisa, Pavonis, and Ascraeus Montes.

ARISA MONS FLOW UNIT 4: Occurs on flanks and lower western and northern slopes of Arisa Mons. Similar to Tharsis Montes flows (unit Aam1) but more subdued and lobate tongues broader; minor faulting. Albedo variable, low to high. Very high resolution frames show surface ridges sub-parallel to lobate frontal scarps. Crater density range, N = 660-840. *Interpretation:* Relatively young lava flows extruded from flanks and possibly central area of Arisa Mons.

ALBA PATERA FLOW UNIT 3: Occurs around and within central caldera. Flows bury and partly bury many ring and radial structures but are cut by others. Channels with levees common along crests of narrow elongate flows. Crater density range, N = 675-1000. *Interpretation:* Youngest lava flows erupted from caldera and fissures during period of intense faulting around central part of Alba Patera.

CERATUNUS FOSSAE FLOWS: Extend southwesterly from highly fractured and faulted region of Ceratunus Fossae. Flows cover or partly cover large areas within fracture zones and overlie Alba Patera flow unit 2 (Ahp2) but are overlapped by Olympus plains and Tharsis Montes flows. Crater density range, N = 850-1150. *Interpretation:* Lava flows extruded from fissures of Ceratunus Fossae and probably other fissure systems northwest of Pavonis Mons.

ARISA MONS FLOW UNIT 3: Embays highland terrain along and basal slopes of Arisa Mons and overlaps Ceratunus Fossae. Lobate flows and ridges less prominent than faults than those along Claritas Fossae. Crater density range, N = 1300-1500. *Interpretation:* Lava flows of intermediate age extruded from fissures in flanks of Arisa Mons.

ARISA MONS FLOW UNIT 2: Occurs on south and southeast flanks of Arisa Mons. Embays highlands and is partly overlapped by unit above and by Tharsis Montes flows. Crater density range, N = 1750-630. *Interpretation:* Lava flows of intermediate age extruded from fissures in flanks of Arisa Mons.

ARISA MONS FLOW UNIT 1: Occurs as relatively small narrow patch on southwest side of Arisa Mons; embays highland terra. Crater density range, N = 1750-1970. *Interpretation:* Oldest exposed lava flows from Arisa Mons.

OLDER FLOWS
VOLCANIC MATERIAL, UNDIVIDED: Forms central promontories of large and small low-relief volcanoes in Tharsis region. Stratigraphic relations indicate a wide range in age for the group. Some small volcanoes may be relatively young, whereas moderate and larger sizes are overlain in places by several flow units. Crater density determinations are not reliable because small volcanic craters and pits occur on the host structures. *Interpretation:* Shield volcanoes formed by the accumulation of lava flows around a central vent throughout extended periods of volcanism in the Tharsis region.

OLYMPUS MONS AUREOLE UNITS 5-1: Occur as a series of large overlapping semicircular lobes surrounding Olympus Mons. Described and interpreted by Morris and Dwornik (1978) as lava flows extruded prior to construction of Olympus Mons. Flows are transected by numerous grooves; therefore, relative ages based on normal crater-retention rates are unreliable. Considered as a group they are older than Arisa Mons flow unit 4 but younger than Alba Patera flow unit 1. Their stratigraphic placement among other flows is based partly on morphology and is somewhat arbitrary. Recent studies by Morris (oral communication, 1980) support an ash-flow origin for the aureoles.

SYRIA PLANUM FLOW UNIT 2: Occurs around depression in crestral area of Syria Planum. Long narrow flows and sheet type flows common; few channels. Partly covers fault systems associated with Claritas Fossae. Transsected in places by troughs of Nectis Labrynthus. Gradational with Syria Planum flow unit 1. Crater density range, N = 1800-2400. *Interpretation:* Lava flows younger than fault systems associated with Claritas Fossae but older than fractures of Nectis Labrynthus. Flows may have been extruded from caldera near present summit of Syria Planum.

ALBA PATERA FLOW UNIT 2: Extends more than 1000 km in places from younger flows (unit Ahp3) cover central part of Alba Patera. Sheet type flows common; some channels with levees. Transsected by many radial and ring fractures. Embays terra of Achernis and Ceratunus Fossae. Gradational with Alba Patera flow unit 1. Same relative age as unit above. Crater density range, N = 1800-2400. *Interpretation:* Volcanic flows from early eruptions of Alba Patera.

SYRIA PLANUM FLOW UNIT 1: Occurs around base of Syria Planum. More highly fractured and flow lobes more subdued but similar in appearance to Syria Planum flow unit 2; boundary only approximately localized. Crater density range, N = 2400-3200. *Interpretation:* Older flows from flank and crestral areas of Syria Planum.

ALBA PATERA FLOW UNIT 1: Similar to Alba Patera flow unit 2 but flow lobes here defined. Boundary with plains materials in northern lowlands not distinct. Embays terra of Achernis Fossae but is overlapped by Olympus Mons aureole unit 3. Originates primarily at Alba Patera where it is cut by peripheral fissure and fault systems, but includes localized, coeval flows apparently extruded radially from the center of a concentric volcano-tectonic feature at 28° N, 126° W. About same relative age as unit above. Crater density range, N = 2400-3200. *Interpretation:* Earliest exposed lava flows from Alba Patera and possibly from other northern eruptions.

SLIDE, CHANNEL, AND FLOOD-PLAIN MATERIALS
SLIDE MATERIAL, UNDIVIDED: Occurs on northwest slopes of Olympus Mons, Arisa Mons, Pavonis Mons, and Ascraeus Montes. Within Nectis Labrynthus. Forms large thin lobes having many small volcanic craters. Crater density range, N = 1300-1500. *Interpretation:* Lava flows extruded from fissures in flanks of Arisa Mons and possibly from other northern eruptions.

CHANNEL AND FLOOD-PLAIN DEPOSITS: Cover floor of Mangala Vallis and adjacent floodplains; in places textural characteristics of substrate visible at high-resolution images. *Interpretation:* This mantle of windblown material.

PLAIN MATERIAL, UNDIVIDED: Forms broad low hills north and west of Mangala Vallis; smooth rolling surfaces, situated in places along margins. Encroached and partly buried adjacent units and landforms. In other areas occurs as broad, relatively level plains that appear rough and striated on moderate-resolution images. Crater density range, N = 850-1150. *Interpretation:* Aluvial deposits of Mangala Vallis and its tributary channels, and restricted occurrences elsewhere.

BASEMENT ROCKS
TERRA MATERIAL, UNDIVIDED: Occurs as both rough and smooth-fractured areas of relatively deformed terrain throughout Tharsis region; forms semicircular ring of Achernis Fossae. Exposed by all other units. Generally highly cratered, but apparent density of craters varies with degree of fracturing. *Interpretation:* Rough surfaces represent ancient volcanic material consisting of volcanic flows and ejecta debris from unknown sources. Smooth surfaces probably mantled by older flows before fracturing formed. Around Achernis Fossae represents rim material of old caldera predating Alba Patera.

NOTE: INDIVIDUAL MAPS MAY NOT SHOW ALL SYMBOLS

Contact: Queried where location uncertain.
Unit outline: Line marks crest; hachures extend down lobate frontal scarp. Dashed where mantled by slide material. Represents contact in place.
Fault or narrow graben: Bar and ball on down-slope side. Dashed where partly buried.
Streamlines: Indicate flow directions where individual units not mapped.
Small channel: Volcanic or fluvial origin.

Escarpment around Olympus Mons: Line marks crest; dashed where buried. Represents contact in places.
Scarp: Line at base, barb points downslope. Represents contact in places.
Ridge crest: Represents contact in places.
Trough:

Caldera or rimless depression:
Impact crater rimless: Represents contact in places.
Impact crater larger than 10 km diameter: Relative age not determined but younger than map unit on which superimposed.
Impact crater larger than 10 km diameter: Older than surrounding map unit.
Impact crater outline on terra material:

MAP SHOWING LAVA FLOWS IN THE SOUTHEAST PART OF THE PHOENICIS LACUS QUADRANGLE OF MARS

By
D.H. Scott
1981

For sale by Branch of Distribution, U.S. Geological Survey,
1200 South East Street, Arlington, VA 22202, and Branch of Distribution,
U.S. Geological Survey, Box 2586, Federal Center, Denver, CO 80225